

The Relativistic Quantum Stationary Hamilton Jacobi Equation for Particle with Spin $1/2$.

T. Djama*

October 29, 2003

Abstract

For one dimensional motions, we derive from the Dirac Spinors Equation (DSE) the Quantum Stationary Hamilton-Jacobi Equation for particles with spin $1/2$. Then, We give its solution. We demonstrate that the $QSHJES_{\frac{1}{2}}$ have two explicit forms, which represent the two possible projection of the Spin $1/2$.

PACS: 03.65.Bz; 03.65.Ca

Key words: relativistic quantum Hamilton-Jacobi equation, spin, conjugate momentum.

*E-mail: djam_touf@yahoo.fr

1- Introduction

During about three years, a new deterministic approach of quantum mechanics was arising from the works of Djama and Bouda [1, 2, 3, 4]. It consist on the introduction of a quantum Lagrangian with which and basing on the Quantum Stationary Hamilton-Jacobi Equation (QSHJE), they derived the Quantum Newton's Law in one dimension [1]. They also plot the quantum trajectories for different potentials [2]. After, Djama has generalize this formalism to the relativistic cases [5, 6]. First, he took up the relativistic QSHJE (RQSHJE) already introduced by Faraggi and Matone [7].

$$\frac{1}{2m_0} \left(\frac{\partial S_0}{\partial x} \right)^2 - \frac{\hbar^2}{4m_0} \left[\frac{3}{2} \left(\frac{\partial S_0}{\partial x} \right)^{-2} \left(\frac{\partial^2 S_0}{\partial x^2} \right)^2 - \left(\frac{\partial S_0}{\partial x} \right)^{-1} \left(\frac{\partial^3 S_0}{\partial x^3} \right) \right] + \frac{1}{2m_0 c^2} [m_0^2 c^4 - (E - V)^2] = 0, \quad (1)$$

and introduced the relativistic quantum Lagrangian written as [5]

$$L = -m_0 c^2 \sqrt{1 - f(x) \frac{\dot{x}^2}{c^2}} - V(x). \quad (2)$$

From this Lagrangian, and using the least action principle, he deduce the expression of the conjugate momentum in function of the particle's velocity [5]

$$\dot{x} \frac{\partial S_0}{\partial x} = E - V(x) - \frac{m_0^2 c^4}{E - V(x)}, \quad (3)$$

from which he derived the Relativistic Quantum Newton's Law

$$\begin{aligned} & [(E - V)^2 - m_0^2 c^4]^2 + \frac{\dot{x}^2}{c^2} (E - V)^2 [(E - V)^2 - m_0^2 c^4] + \frac{\hbar^2}{2} \left[\frac{3}{2} \left(\frac{\ddot{x}}{\dot{x}} \right)^2 - \frac{\ddot{x}}{\dot{x}} \right] \cdot \\ & (E - V)^2 - \frac{\hbar^2}{2} \left(\ddot{x} \frac{dV}{dx} + \dot{x}^2 \frac{d^2 V}{dx^2} \right) \left[\frac{(E - V)^2 + m_0^2 c^4}{(E - V)^2 - m_0^2 c^4} \right] (E - V)^2 - \frac{3\hbar^2}{4} \cdot \\ & \left(\dot{x} \frac{dV}{dx} \right)^2 \left[\frac{(E - V)^2 + m_0^2 c^4}{(E - V)^2 - m_0^2 c^4} \right]^2 - \hbar^2 \left(\dot{x} \frac{dV}{dx} \right)^2 \frac{m_0^2 c^4}{(E - V)^2 - m_0^2 c^4} = 0. \quad (4) \end{aligned}$$

In a last paper [6], Djama had drawn the trajectories of a relativistic spinless particles for many potentials, and established the existence of nodes throw which all possible trajectories pass, even the purely relativistic one [6]. It is useful to stress that those results are found in one dimension.

It is clear that the introduction of quantum and relativistic quantum trajectories in the concept of a deterministic approach of quantum and relativistic quantum mechanics is a great step to elaborate a deterministic theory, but it is not sufficient. The generalization to spinning particle case and three dimension problems must be investigated.

The object of this paper is to investigate -in one dimension- the Law of motion of particles with spin 1/2. With this aim, we derive in Sec. 2 the RQSHJE

for Spin 1/2 (RQSHJES_{1/2}) from the Dirac Spinors Equation (DSE) written in one dimension. We demonstrate that the RQSHJES_{1/2} is composed by two components, each one correspond to the projection of the Spin 1/2 ($m_s = +1/2$ and $m_s = -1/2$). Then, in Sec. 3, we propose a Double solution of the RQSHJES_{1/2} corresponding to $m_s = +1/2$ and $m_s = -1/2$. Finally, in Sec. 4, we discuss our results.

2- Quantum Stationary Hamilton-Jacobi Equation for a particle with Spin 1/2

In the beginning, the quantum systems are described by the Schrödinger equation

$$-\frac{\hbar^2}{2m} \frac{d^2\psi}{dx^2} + V(x) = E . \quad (5)$$

But later, after the discovery of the spin by Goodsmith [8] and Pauli [8], and for studying the spinning particle's behaviour, Dirac proposed his famous Spinors equation, written with Pauli representation as [9]

$$i \hbar c \vec{\alpha} \cdot \vec{\nabla}(\Psi) = [E - V(\vec{r}) - \beta m_0 c^2] \quad (6)$$

where $\alpha_i, i = (x, y, z)$ and β are the Dirac matrices

$$\alpha_i = \begin{pmatrix} 0 & \sigma_i \\ \sigma_i & 0 \end{pmatrix}; \quad \beta = \begin{pmatrix} I & 0 \\ 0 & -I \end{pmatrix}, \quad (7)$$

for which σ_i are the Pauli matrices (Eq. (9)), and $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ is the identity matrix.

The Dirac equation is written, in one dimension, in the Appendix of Ref. [10] as

$$-i \hbar c \sigma_x \frac{d\psi}{dx} = (E - V(x) - \sigma_z m_0 c^2) \psi \quad (8)$$

where

$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}; \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}; \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (9)$$

are the Pauli matrix. $\psi = \begin{pmatrix} \theta \\ \phi \end{pmatrix}$ is a the matrix of the wave functions θ and ϕ .

Eq. (7) is a matrix equation of two components describing the state of particles of half spin with energy E moving in one dimension under the action of the potential $V(x)$. It contains two scalar equations which can be deduced after decomposing

$$-i \hbar c \frac{d\phi}{dx} = (E - V(x) - m_0 c^2) \theta(x) , \quad (10)$$

$$-i \hbar c \frac{d\theta}{dx} = (E - V(x) + m_0 c^2) \phi(x) . \quad (11)$$

Taking the expression of θ from Eq. (10) and replacing it into Eq. (11), then, taking the expression of ϕ from Eq. (11) and replacing it into Eq. (10), we find

$$\hbar^2 c^2 \frac{d^2 \theta}{dx^2} + \hbar^2 c^2 \frac{dV/dx}{E - V + m_0 c^2} \frac{d\theta}{dx} + [(E - V)^2 - m_0^2 c^4] \theta(x) = 0, \quad (12)$$

$$\hbar^2 c^2 \frac{d^2 \phi}{dx^2} + \hbar^2 c^2 \frac{dV/dx}{E - V - m_0 c^2} \frac{d\phi}{dx} + [(E - V)^2 - m_0^2 c^4] \phi(x) = 0. \quad (13)$$

Now, the next step is to establish the QSHJES $_{\frac{1}{2}}$. In this order, let us write the two wave functions as

$$\theta(x) = A(x) \left(\alpha_+ e^{\frac{i}{\hbar} S_0} + \alpha_- e^{-\frac{i}{\hbar} S_0} \right), \quad (14)$$

$$\phi(x) = B(x) \left(\beta_+ e^{\frac{i}{\hbar} Z_0} + \beta_- e^{-\frac{i}{\hbar} Z_0} \right), \quad (15)$$

where $A(x)$, $B(x)$, $S_0(x)$ and $Z_0(x)$ are real functions of x . α_+ , α_- , β_+ and β_- are real constants.

Replacing Eqs. (14) and (15) into Eqs.(12) and (13) respectively and decomposing it, we find

$$\frac{\hbar^2 c^2}{A} \frac{d^2 A}{dx^2} - c^2 \left(\frac{dS_0}{dx} \right)^2 + \hbar^2 c^2 \frac{dV/dx}{E - V + m_0 c^2} \frac{1}{A} \frac{dA}{dx} + [(E - V)^2 - m_0^2 c^4] = 0, \quad (16)$$

$$\frac{\hbar^2 c^2}{B} \frac{d^2 B}{dx^2} - c^2 \left(\frac{dZ_0}{dx} \right)^2 + \hbar^2 c^2 \frac{dV/dx}{E - V - m_0 c^2} \frac{1}{B} \frac{dB}{dx} + [(E - V)^2 - m_0^2 c^4] = 0, \quad (17)$$

and

$$A \frac{d^2 S_0}{dx^2} + 2 \frac{dA}{dx} \frac{dS_0}{dx} + \frac{dV/dx}{E - V + m_0 c^2} A \frac{dS_0}{dx} = 0, \quad (18)$$

$$B \frac{d^2 Z_0}{dx^2} + 2 \frac{dB}{dx} \frac{dZ_0}{dx} + \frac{dV/dx}{E - V - m_0 c^2} B \frac{dZ_0}{dx} = 0. \quad (19)$$

We can show easily that Eqs. (18) and (19) can be integrated to give

$$A(x) = k_1 (E - V(x) + m_0 c^2)^{\frac{1}{2}} \left(\frac{dS_0}{dx} \right)^{-\frac{1}{2}}, \quad (20)$$

$$B(x) = k_2 (E - V(x) - m_0 c^2)^{\frac{1}{2}} \left(\frac{dZ_0}{dx} \right)^{-\frac{1}{2}}, \quad (21)$$

where k_1 and k_2 are two real constants. Replacing Eqs. (20) and (21) into Eqs. (16) and (17), we get

$$\begin{aligned} & \frac{1}{2m_0} \left(\frac{dS_0}{dx} \right)^2 - \frac{\hbar^2}{4m_0} \{S_0, x\} + \frac{\hbar^2}{2m_0} (E - V + m_0 c^2)^{\frac{1}{2}} \\ & \cdot \frac{d^2}{dx^2} \left[(E - V + m_0 c^2)^{-\frac{1}{2}} \right] + \frac{1}{2m_0 c^2} [m_0^2 c^4 - (E - V)^2] = 0, \end{aligned} \quad (22)$$

$$\begin{aligned} & \frac{1}{2m_0} \left(\frac{dZ_0}{dx} \right)^2 - \frac{\hbar^2}{4m_0} \{Z_0, x\} + \frac{\hbar^2}{2m_0} (E - V - m_0 c^2)^{\frac{1}{2}} \\ & \cdot \frac{d^2}{dx^2} \left[(E - V - m_0 c^2)^{-\frac{1}{2}} \right] + \frac{1}{2m_0 c^2} [m_0^2 c^4 - (E - V)^2] = 0, \end{aligned} \quad (23)$$

where

$$\{f(x), x\} = \left[\frac{3}{2} \left(\frac{df}{dx} \right)^{-2} \left(\frac{d^2f}{dx^2} \right)^2 - \left(\frac{df}{dx} \right)^{-1} \left(\frac{d^3f}{dx^3} \right) \right]$$

represents the schwarzian derivative of $f(x)$ with respect to x . Eqs. (22) and (23) represent the two Relativistic Quantum Stationary Hamilton Jacobi Equations for Spinning particle ($s = \frac{1}{2}$) (QSHJES $_{\frac{1}{2}}$). One of these equations correspond to the projection $m_s = +\frac{1}{2}$ of the spin, when the other correspond to the projection $m_s = -\frac{1}{2}$. It follows that the reduced actions S_0 and Z_0 correspond to the two projections of the spin.

We can connect the wave functions θ and ϕ to the reduced actions as follows

$$\psi = \begin{pmatrix} \theta \\ \phi \end{pmatrix} = \begin{pmatrix} k_1(E - V(x) + m_0c^2)^{\frac{1}{2}} \left(\frac{dS_0}{dx} \right)^{-\frac{1}{2}} \left(\alpha_+ e^{\frac{i}{\hbar} S_0} + \alpha_- e^{-\frac{i}{\hbar} S_0} \right) \\ k_2(E - V(x) - m_0c^2)^{\frac{1}{2}} \left(\frac{dZ_0}{dx} \right)^{-\frac{1}{2}} \left(\beta_+ e^{\frac{i}{\hbar} Z_0} + \beta_- e^{-\frac{i}{\hbar} Z_0} \right) \end{pmatrix} \quad (24)$$

Now, let us discuss Eqs. (22) and (23). First, remark that, compared to the RQSHJE (Eq. (1)), Eq. (22) and (23) contains additional terms

$$term_1 = \frac{\hbar^2}{2m_0} (E - V + m_0c^2)^{\frac{1}{2}} \frac{d^2}{dx^2} \left[(E - V + m_0c^2)^{-\frac{1}{2}} \right] \quad (25)$$

for Eq. (22), and

$$term_1 = \frac{\hbar^2}{2m_0} (E - V - m_0c^2)^{\frac{1}{2}} \frac{d^2}{dx^2} \left[(E - V - m_0c^2)^{-\frac{1}{2}} \right] \quad (26)$$

for Eq. (23). Both $term_1$ and $term_2$ are proportional to $\hbar^2$ which means that at the classical limit ($\hbar \rightarrow 0$) they vanish and Eqs. (22) and (23) reduce to the Classical Stationary Hamilton-Jacobi Equation (CSHJE). Secondly, these two terms vanish in the case of a constant potential. So, Eqs. (22) and (23) reduce to the QSHJE. More than this, the motion of a spinning particles are not affected by the constant potentials but only by potentials with a not vanishing gradient with respect to x .

By this, we deduce that $term_1$ and $term_2$ are directly related to the spinning behaviour of quantum particles in one dimension.

3- The solutions of the RQSHJES $_{\frac{1}{2}}$

Now, let us investigate the form of the reduced actions S_0 and Z_0 solutions of the RQSHJES $_{\frac{1}{2}}$ (Eqs. (22) and (23)). We propose as solutions of Eqs. (22) and (23) the following functions

$$S_0 = \hbar \arctan \left(a \frac{\theta_1(x)}{\theta_2(x)} + b \right), \quad (27)$$

$$Z_0 = \hbar \arctan \left(d \frac{\phi_1(x)}{\phi_2(x)} + e \right), \quad (28)$$

where a, b, d and e are real constants. θ_1 and θ_2 are two real and independent solutions of Eq. (12). ϕ_1 and ϕ_2 are two real independent solutions of Eq. (13).

Expressions (27) and (28) are analogous to the expression of the reduced action, solution of the RQSHJE and the QSHJE [1, 5, 7, 11].

Now, we demonstrate our proposition. First, remark that Eqs. (22) and (23) have an analogous form. To write Eq. (23) we can replace S_0 by Z_0 and $(E - V + m_0 c^2)$ by $(E - V - m_0 c^2)$. Eqs. (27) and (28) are also analogous. For this reason, in what follows, we exhibit demonstration only for S_0 and Eq. (22) (for Z_0 and Eq. (23) the demonstration can be done with the same manner). In this order, we introduce the function

$$\theta = a \theta_1 + b \theta_2 . \quad (29)$$

As θ_1 and θ_2 are solutions of Eq. (12), θ is also solution of it. S_0 takes the form

$$S_0 = \hbar \arctan \left(a \frac{\theta(x)}{\theta_2(x)} \right) . \quad (30)$$

The wronskian W of θ and θ_2 is deduced from Eq. (12)

$$W = \theta \frac{d\theta_2}{dx} - \theta_2 \frac{d\theta}{dx} = \alpha (E - V(x) + m_0 c^2) , \quad (31)$$

where α is a real constant. Using this expression of the wronskian and replacing Eq. (30) into Eq. (22), after calculating the schwarzian derivative of S_0 with respect to x , one find

$$\begin{aligned} & \frac{\hbar^2}{2m_0} \frac{W^2 - (\theta \frac{d\theta_2}{dx} - \theta_2 \frac{d\theta}{dx})^2}{(\theta^2 + \theta_2^2)^2} - \frac{\hbar^2}{2m_0} (E - V + m_0 c^2)^{\frac{1}{2}} \cdot \\ & \cdot \frac{d^2}{dx^2} \left[(E - V + m_0 c^2)^{-\frac{1}{2}} \right] + \frac{\hbar^2}{2m_0} (E - V + m_0 c^2)^{\frac{1}{2}} \cdot \\ & \cdot \frac{d^2}{dx^2} \left[(E - V + m_0 c^2)^{-\frac{1}{2}} \right] - \frac{\hbar^2}{2m_0} \left[\frac{dV/dx}{E - V + m_0 c^2} \cdot \right. \\ & \cdot \left. \frac{(\theta \frac{d\theta_2}{dx} - \theta_2 \frac{d\theta}{dx})}{\theta^2 + \theta_2^2} + \frac{(\theta \frac{d^2\theta_2}{dx^2} + \theta_2 \frac{d^2\theta}{dx^2})}{\theta^2 + \theta_2^2} \right] \\ & + \frac{1}{2m_0 c^2} [m_0^2 c^4 - (E - V)^2] = 0 , \end{aligned} \quad (32)$$

which reduces to

$$\begin{aligned} & \theta \left\{ \hbar^2 c^2 \frac{d^2\theta}{dx^2} + \hbar^2 c^2 \frac{dV/dx}{E - V + m_0 c^2} \frac{d\theta}{dx} + [(E - V)^2 - m_0^2 c^4] \theta \right\} \\ & + \theta_2 \left\{ \hbar^2 c^2 \frac{d^2\theta_2}{dx^2} + \hbar^2 c^2 \frac{dV/dx}{E - V + m_0 c^2} \frac{d\theta_2}{dx} + [(E - V)^2 - m_0^2 c^4] \theta_2 \right\} = 0 . \end{aligned} \quad (33)$$

As θ and θ_2 are two solutions of Eq. (12), Eq. (33) is automatically satisfied. Then, expression given by Eq. (27) is the solution of the RQSHJE $_{\frac{1}{2}}$ given by Eq. (22). We note that, with same manner, we can demonstrate that expression given by Eq. (28) is the solution of the RQSHJE $_{\frac{1}{2}}$ given by Eq. (23).

4- conclusion

In this paper, we have present two new results. The first one is the establishment of the two Relativistic Quantum Stationary Hamilton-Jacobi Equations for a half spinning particle ($s = \frac{1}{2}$) (QSHJES $_{\frac{1}{2}}$). The second result is the resolution of the RQSHJE $_{\frac{1}{2}}$.

The RQSHJE $_{\frac{1}{2}}$ is composed of two equations (Eq. (22) and (23)). Each one represent a projection of the spin ($m_s = \frac{1}{2}$ and $m_s = -\frac{1}{2}$).

both RQSHJE $_{\frac{1}{2}}$ reduce to the classical Hamilton-Jacobi equation when we take the classical limit ($\hbar \rightarrow 0$).

for a free particle the spinning terms (Eqs. (25) and (26)) vanish and both RQSHJE $_{\frac{1}{2}}$ reduce to the RQSHJE. So, for the free particle, we can not distinguish between two projection of spin until it pass throw a non vanishing gradient potential.

Another interesting remark is the non relativistic case. the question is: Can we deduce, from the RQSHJE $_{\frac{1}{2}}$, the spinning behaviour of a particle for a purely quantum case? The answer is: Yes. Let us review the RQSHJE $_{\frac{1}{2}}$ (Eqs. (22) and (23)). For the non relativistic limit, $T \ll m_0 c^2$ (where $T = E - V(x) - m_0 c^2$ is the kinetic energy of the particle), Eq. (22) reduces to

$$\frac{1}{2m_0} \left(\frac{dS_0}{dx} \right)^2 - \frac{\hbar^2}{4m_0} \{S_0, x\} + V(x) - E' = 0 \quad (34)$$

which is the ordinary QSHJE. E' is the quantum energy without rest energy $m_0 c^2$.

In the other hand side, Eq. (23) reduces to

$$\begin{aligned} \frac{1}{2m_0} \left(\frac{dZ_0}{dx} \right)^2 - \frac{\hbar^2}{4m_0} \{Z_0, x\} + \frac{\hbar^2}{2m_0} (E' - V)^{\frac{1}{2}} . \\ \cdot \frac{d^2}{dx^2} \left[(E' - V)^{-\frac{1}{2}} \right] + V(x) - E' = 0 , \end{aligned} \quad (35)$$

which is different from the ordinary QSHJE. So, it is clear that Eq. (35) describe one of the two projections of the spin. Thus there is two QSHJES $_{\frac{1}{2}}$ to describe the spinning particle with $s = \frac{1}{2}$ in the purely quantum cases. This point will be more investigated in a next papers.

REFERENCES

1. A. Bouda and T. Djama, *Phys. Lett. A* 285 (2001) 27-33; quant-ph/0103071.
2. A. Bouda and T. Djama, ; *Physica scripta* 66 (2002) 97-104; quant-ph/0108022.
3. A. Bouda and T. Djama, *Phys. Lett. A* 296 (2002) 312-316; quant-ph/0206149.
4. E. R. Floyd, *Phys. Lett. A* 296 (2002) 307-311; quant-ph/0206114.
5. T. Djama, "Relativistic Quantum Newton's Law and photon Trajectories"; quant-ph/0111121.
6. T. Djama, "Nodes in the Relativistic Quantum Trajectories and Photon's Trajectories ;quant-ph/0201003.
7. A. E. Faraggi and M. Matone, *Int. J. Mod. Phys. A* 15, 1869 (2000); hep-th/9809127.
8. E. Chpolski, *Physique Atomique*, Tome II, Edition Mir.
9. Any bibliography on the quantum field theory. See also:
Guang-jiong NI and Tsao CHANG, "Is Neutrino a Superluminal Particle?"; hep-ph/0103051.
Zhi-Jian Li , J. Q. Liang , D. H. Kobe, "Larmor precession and barrier tunneling time of a neutral spinning particle"; quant-ph/0109001
10. H. Nitta, T. Kudo, and H. Minowa, "Motion of a wave packet in the Klein paradox," *Am. J. Phys.* 67, 966-971 (1999).
11. E. R. Floyd, *Phys. Rev. D* 34, 3246 (1986).